



12

EUROPEAN PATENT SPECIFICATION

45 Date of publication of patent specification :
19.01.94 Bulletin 94/03

51 Int. Cl.⁵ : **C07C 401/00, A61K 31/59**

21 Application number : **90302521.1**

22 Date of filing : **09.03.90**

54 **19-Nor vitamin D compounds.**

30 Priority : **09.03.89 US 321030**
16.02.90 US 481354

43 Date of publication of application :
12.09.90 Bulletin 90/37

45 Publication of the grant of the patent :
19.01.94 Bulletin 94/03

84 Designated Contracting States :
AT BE CH DE DK ES FR GB GR IT LI LU NL SE

66 References cited :
EP-A- 0 250 755
WO-A-85/03300
WO-A-86/02649
US-A- 4 448 726
JOURNAL OF THE CHEMICAL SOCIETY PER-
KIN TRANSACTIONS I, vol. 6, 1978, pages
590-595, London, GB; B. LYTHGOE et al.:
"Calciferol and its relatives. Part 22. A direct
total synthesis of vitamin D2 and vitamin
D3*2"

73 Proprietor : **WISCONSIN ALUMNI RESEARCH**
FOUNDATION
614 North Walnut Street
Madison, WI 53705 (US)

72 Inventor : **DeLuca, Hector F.**
1809 Highway BB Deerfield
Wisconsin 53531 (US)
Inventor : **Schnoes, Heinrich K.**
1806 Summit Avenue
Madison Wisconsin 53705 (US)
Inventor : **Periman, Kato L.**
1 Chippewa Court
Madison Wisconsin 53711 (US)
Inventor : **Sicinski, Rafal R.**
University of Warsaw Dept. of Chemistry
02-093 Warsaw Pasteura 1 (PL)
Inventor : **Prahl, Jean Martin**
7 Georgetown Court
Madison Wisconsin 53919 (US)

74 Representative : **Ellis-Jones, Patrick George**
Armine et al
J.A. KEMP & CO. 14 South Square Gray's Inn
London WC1R 5LX (GB)

EP 0 387 077 B1

Note : Within nine months from the publication of the mention of the grant of the European patent, any person may give notice to the European Patent Office of opposition to the European patent granted. Notice of opposition shall be filed in a written reasoned statement. It shall not be deemed to have been filed until the opposition fee has been paid (Art. 99(1) European patent convention).

Description

This invention was made with United States government support awarded by the Department of Health and Human Services (NIH), Grant number: DK-14881. The United States Government has certain rights in this invention.


This invention relates to biologically active vitamin D compounds. More specifically, the invention relates to 19-nor-analogs of 1 α -hydroxylated vitamin D compounds and to a general process for their preparation.

Background


The 1 α -hydroxylated metabolites of vitamin D – most importantly 1 α ,25-dihydroxyvitamin D₃ (See US-A-3697559) and 1 α ,25-dihydroxyvitamin D₂ – are known as highly potent regulators of calcium homeostasis in animals and humans, and more recently their activity in cellular differentiation has also been established. As a consequence, many structural analogs of these metabolites, such as compounds with different side chain structures, different hydroxylation patterns, or different stereochemistry, have been prepared and tested. Important examples of such analogs are 1 α -hydroxyvitamin D₃, 1 α -hydroxyvitamin D₂, various side chain fluorinated derivatives of 1 α ,25-dihydroxyvitamin D₃, and side chain homologated analogs. Several of these known compounds exhibit highly potent activity *in vivo* or *in vitro*, and possess advantageous activity profiles and thus are in use, or have been proposed for use, in the treatment of a variety of diseases such as renal osteocystrophy, vitamin D-resistant rickets, osteoporosis, psoriasis, and certain malignancies.

Disclosure and Description of the Invention

A class of 1 α -hydroxylated vitamin D compounds not known heretofore are the 19-nor-analogs, i.e. compounds in which the ring A exocyclic methylene group (carbon 19) typical of all vitamin D system has been removed and replaced by two hydrogen atoms. Structurally these novel analogs are characterized by the general formula I shown below:




where X¹ and X² are each selected from the group consisting of hydrogen and acyl, and where the group R represents any of the typical side chains known for vitamin D type compounds. Thus, R may be an alkyl, hydrogen, hydroxyalkyl or fluoroalkyl group, or R may represent the following side chain:



wherein R¹ represents hydrogen, hydroxy or O-acyl, R² and R³ are each selected from the group consisting of alkyl, hydroxyalkyl and fluoroalkyl, or, when taken together represent the group – (CH₂)_m – where m is an integer having a value of from 2 to 5, R⁴ is selected from the group consisting of hydrogen, hydroxy, fluorine, O-


acyl, alkyl, hydroxyalkyl and fluoroalkyl, R^5 is selected from the group consisting of hydrogen, fluorine, alkyl, hydroxyalkyl and fluoroalkyl, or, R^4 and R^6 taken together represent double-bonded oxygen, R^6 and R^7 are each selected from the group consisting of hydrogen, hydroxy, O-acyl, fluorine and alkyl, or, R^6 and R^7 taken together form a carbon-carbon double bond, and wherein n is an integer having a value of from 1 to 5, and wherein the carbon at any one of positions 20, 22, or 23 in the side chain may be replaced by an O, S, or N atom.

Specific important examples of side chains are the structures represented by formulas (a), (b), (c), (d) and (e) below; i.e. the side chain as it occurs in 25-hydroxyvitamin D₃ (a); vitamin D₃ (b); 25-hydroxyvitamin D₂ (c); vitamin D₂ (d); and the C-24-epimer of 25-hydroxyvitamin D₂ (e).




In this specification and the claims, the term 'alkyl' signifies an alkyl radical of 1 to 5 carbons in all isomeric forms, such as methyl, ethyl, propyl, isopropyl, butyl, isobutyl, pentyl, etc., and the terms 'hydroxyalkyl' and 'fluoroalkyl' refer to such an alkyl radical substituted by one or more hydroxy or fluoro groups respectively, and the term 'acyl' means an aliphatic acyl group of 1 to 5 carbons, such as formyl, acetyl, propionyl, etc. or an aromatic acyl group such as benzoyl, nitrobenzoyl or halobenzoyl. The term 'aryl' signifies a phenyl-, or an alkyl-, nitro- or halo-substituted phenyl group.

The preparation of 1 α -hydroxy-19-nor-vitamin D compounds having the basic structure shown above can be accomplished by a common general method, using known vitamin D compounds as starting materials. Suitable starting materials are, for example, the vitamin D compounds of the general structure II:




where R is any of the side chains as defined above. These vitamin D starting materials are known compounds, or compounds that can be prepared by known methods.

Using the procedure of DeLuca *et al.* (U.S. Patent 4,195,027), the starting material can be converted to the corresponding 1 α -hydroxy-3,5-cyclovitamin D derivative, having the general structure III below, where X represents hydrogen and Q represents an alkyl, preferably methyl:



So as to preclude undesired reaction of the 1 α -hydroxy group in subsequent steps, the hydroxy group is converted to the corresponding acyl derivative, i.e. the compound III shown above, where X represents an acyl group, using standard acylation procedures, such as treatment with an acyl anhydride or acyl halide in pyridine at room temperature or slightly elevated temperature (30-70°C). It should be understood also that whereas the process of this invention is illustrated here with acyl protection of hydroxy functions, alternative standard hydroxy-protecting groups can also be used, such as, for example, alkylsilyl or alkoxyalkyl groups. Such protecting groups are well-known in the art (e.g. trimethylsilyl, triethylsilyl, t-butyldimethylsilyl, or tetrahydrofuranyl, methoxymethyl), and their use is considered a routine modification of experimental detail within the scope of the process of this invention.


The derivative as obtained above can then be reacted with, say osmium tetroxide, to produce the 10,19-dihydroxy analog, IV (where X is acyl), which is subjected to diol cleavage using sodium metaperiodate or similar vicinal diol cleavage reagents (e.g. lead tetraacetate) to obtain the 10-oxo-intermediate, having the structure V below (where X is acyl):



These two consecutive steps can be carried out according to the procedures given by Paaren *et al.* [J. Org. Chem. 48, 3819 (1983)]. If the side chain unit, R, carries vicinal diols (e.g. 24,25-dihydroxy- or 25,26-dihydroxy, etc.), these, of course, also need to be protected, e.g. via acylation, silylation, or as the isopropylidene derivative prior to the periodate cleavage reactions.


In most cases, the acylation of the 1 α -hydroxy group as mentioned above will simultaneously effect the acylation of side chain hydroxy functions, and these acylation conditions can, of course, be appropriately adjusted (e.g. elevated temperatures, longer reaction times) so as to assure complete protection of side chain vicinal diol groupings.

The next step of the process comprises the reduction of the 10-oxo-group to the corresponding 10-alcohol having the structure VI shown below (where X is acyl and Y represents hydroxy). When X is acyl, this reduction is carried out conveniently in an organic solvent at from, say, 0°C to room temperature, using NaBH₄ or equivalent hydride reducing agents, selective for the reduction of carbonyl groups without cleaving ester functions. Obviously, when X is a hydroxy-protecting group that is stable to reducing agents, any of the other hydride reducing agents (e.g. LiAlH₄, or analogous reagents) may be employed also.



The 10-hydroxy intermediate can then be treated with an alkyl- or arylsulfonylhalide (e.g. methanesulfonylchloride) in a suitable solvent (e.g. pyridine) to obtain the corresponding 10-O-alkyl- or arylsulfonyl derivative (the compound having the structure shown VI above, where Y is alkyl-SO₂O-, or aryl-SO₂O-, and this sulfonate intermediate is then directly reduced, e.g. with lithium aluminum hydride, or the analogous known lithium aluminum alkyl hydride reagents in an ether solvent, at a temperature typically from 0°C to the boiling temperature of the solvent, thereby displacing the sulfonate group and obtaining the 10-deoxy derivative, represented by the structure VI above, where X and Y are both hydrogen. As shown by the above structure, a 1-O-acyl function in the precursor compound V is also cleaved in this reduction step to produce the free 1α-hydroxy function, and any O-acyl protecting group in the side chain would, of course, likewise be reduced to the corresponding free alcohol function, as is well understood in the art. If desired, the hydroxy groups at C-1 (or hydroxy groups in the side chain) can be reprotected by acylation or silylation or other formation to the corresponding acyl, alkylsilyl or alkoxyalkyl derivative, but such protection is not required. Alternative hydroxy-protecting groups, such as alkylsilyl or alkoxyalkyl groups would be retained in this reduction step, but can be removed, as desired, at this or later stages in the process by standard methods known in the art.

The above 1α-hydroxy-10-deoxy cyclovitamin D intermediate is next solvolysed e.g. the presence of a low-molecular weight organic acid, using the conditions of DeLuca *et al.* (U.S. Patents 4,195,027 and 4,260,549). When the solvolysis is carried out in acetic acid, for example, there is obtained a mixture of 1α-hydroxy-19-nor-vitamin D 3-acetate and 1α-hydroxy-19-nor-vitamin D 1-acetate (compounds VII and VIII, below), and the analogous 1- and 3-acylates are produced, when, alternative acids are used for solvolysis.



Direct basic hydrolysis of this mixture under standard conditions then produces the desired 1α-hydroxy-19-nor-vitamin D compounds of structure I above (where X¹ and X² are hydrogen). Alternatively, the above mixture of monoacetates or other acylates may also be separated (e.g. by high pressure liquid chromatography) and the resulting 1-acetate and 3-acetate isomers may be subjected separately to hydrolysis to obtain the same

final product from each, namely the 1α -hydroxy-19-nor-vitamin D compounds of structure I. Also the separated monoacetates of structure VII or VIII or the free 1,3-dihydroxy compound can, of course, be reacylated according to standard procedures with any desired acyl group, so as to produce the product of structure I above, where X^1 and X^2 represent acyl groups which may be the same or different.

Biological Activity of 1α -Hydroxy-19-Nor-Vitamin D Compounds

The novel compounds of this invention exhibit an unexpected pattern of biological activity, namely high potency in promoting the differentiation of malignant cells and little or no activity in calcifying bone tissue. This is illustrated by the biological assay results obtained for $1\alpha,25$ -dihydroxy-19-nor-vitamin D_3 (compounds Ia), which are summarized in Tables 1 and 2, respectively. Table 1 shows a comparison of the activity of the known active metabolite $1\alpha,25$ -dihydroxyvitamin D_3 and the 19-nor analog (Ia) in inducing the differentiation of human leukemia cells (HL-60 cells) in culture to normal cells (monocytes). Differentiation activity was assessed by three standard differentiation assays, abbreviated in Table 1 as NBT (nitroblue tetrazolium reduction), NSE (non-specific esterase activity), and PHAGO (phagocytosis activity). The assays were conducted according to known procedures, as given, for example, by DeLuca *et al.* (U.S. Patent 4,717,721) and Ostrem *et al.*, J. Biol. Chem. **262**, 14164, 1987). For each assay, the differentiation activity of the test compounds is expressed in terms of the percent of HL-60 cells having differentiated to normal cells in response to a given concentration of test compound.

The results summarized in Table 1 clearly show that the new analog, $1\alpha,25$ -dihydroxy-19-nor-vitamin D_3 (Ia) is as potent as $1\alpha,25$ -dihydroxyvitamin D_3 in promoting the differentiation of leukemia cells. Thus in all three assays close to 90% of the cells are induced to differentiate by $1\alpha,25$ -dihydroxyvitamin D_3 at a concentration of 1×10^{-7} molar, and the same degree of differentiation (i.e. 90, 84 and 90%) is achieved by the 19-nor analog (Ia).

Table 1
Differentiation of HL-60 Cells

$1\alpha,25$ -dihydroxyvitamin D_3 (moles/liter)	Z Differentiated Cells (mean \pm SEM)		
	NBT	NSE	PHAGO
1×10^{-7}	86 \pm 2	89 \pm 1	87 \pm 3
1×10^{-8}	60 \pm 2	60 \pm 3	64 \pm 2
1×10^{-9}	33 \pm 2	31 \pm 2	34 \pm 1
 $1\alpha,25$ -Dihydroxy-19-nor- vitamin D_3 , (Ia)			
(moles/liter)			
2×10^{-7}	94 \pm 2	95 \pm 3	94 \pm 2
1×10^{-7}	90 \pm 4	84 \pm 4	90 \pm 4
5×10^{-8}	72 \pm 3	73 \pm 3	74 \pm 3
1×10^{-8}	61 \pm 3	60 \pm 3	56 \pm 1
1×10^{-9}	32 \pm 1	31 \pm 1	33 \pm 1

In contrast to the preceding results, the new 19-nor analog (Ia) exhibits no activity in an assay measuring the calcification of bone, a typical response elicited by vitamin D compounds. Relevant data, representing the

results of an assay comparing the bone calcification activity in rats of 1 α ,25-dihydroxyvitamin D₃ and 1 α ,25-dihydroxy-19-nor-vitamin D₃ (Ia), are summarized in Table 2. This assay was conducted according to the procedure described by Tanaka *et al.*, *Endocrinology* 92, 417 (1973).

The results presented in Table 2 show the expected bone calcification activity of 1 α ,25-dihydroxyvitamin D₃ as reflected by the increase in percent bone ash, and in total ash at all dose levels. In contrast, the 19-nor analog Ia exhibits no activity at all three dose levels, when compared to the vitamin D-deficient (-D) control group.

Table 2.

Calcification Activity

<u>Compound</u>	<u>Amount Administered</u> [*] (pmoles/day/7 days)	<u>% Ash</u> (mean \pm SEM)	<u>Total Ash (mg)</u> (mean \pm SEM)
-D (control)	0	19 \pm 0.8	23 \pm 1.2
1 α ,25-dihydroxy- vitamin D ₃	32.5 65.0 325.0	23 \pm 0.5 26 \pm 0.7 28 \pm 0.9	34 \pm 1.6 36 \pm 1.1 40 \pm 1.9
1 α ,25-dihydroxy-19- nor-vitamin D ₃ (Ia)	32.5 65.0 325.0	22 \pm 0.9 19 \pm 1.5 19 \pm 1.2	28 \pm 1.6 28 \pm 3.4 30 \pm 2.4

* Each assay group comprised 6 rats, receiving the indicated amount of test compound by intraperitoneal injection daily for a period of seven days.

Thus the new 19-nor analog shows a selective activity profile combining high potency in inducing the differentiation of malignant cells with very low or no bone calcification activity. The compounds of this novel structural class, therefore, can be useful as therapeutic agents for the treatment of malignancies. Because the differentiative activity of vitamin D compounds on keratinocytes of skin (Smith *et al.*, *J. Invest. Dermatol.* 88, 709, 1986; Smith *et al.*, *J. Am. Acad. Dermatol.* 19, 516, 1988) is believed to be an indication of successful treatment of psoriasis (Takamoto *et al.*, *Calc. Tissue Int.* 39, 360, 1986), these compounds should prove useful in treating this and other skin disorders characterized by proliferation of undifferentiated skin cells. These compounds should also find use in the suppression of parathyroid tissue, as for example, in cases of secondary hyperparathyroidism found in renal disease (Slatopolsky *et al.*, *J. Clin. Invest.* 74, 2136, 1984).

For treatment purposes, the novel compounds of this invention can be formulated in a solid or liquid vehicle ingestible by, and non toxic to, mammals, for example as solutions in innocuous solvents or as emulsions, suspensions or dispersions in suitable innocuous solvents or carriers, or as pills, tablets or capsules, containing solid carriers according to conventional methods known in the art. Accordingly the present invention provides a pharmaceutical composition which comprises at least one compound of this invention together with a pharmaceutically acceptable excipient. For topical applications or administrations the compounds are advantageously formulated as creams or ointments or similar vehicle suitable for topical applications. Any such formulations may also contain other pharmaceutically-acceptable and non-toxic excipients such as stabilizers, antioxidants, binders, coloring agents or emulsifying or taste-modifying agents.

The compounds are advantageously administered by injection (for parenteral administration), or by intra-

venous infusion of suitable sterile solutions, or in the form of oral doses (for oral administration) via the alimentary canal, or topically in the form of ointments, lotions, or in suitable transdermal patches. For the treatment of malignant diseases, the 19-nor-vitamin D compounds of this invention should be administered to subjects in dosages sufficient to inhibit the proliferation of malignant cells and induce their differentiation into normal monocyte-macrophages. Similarly, for the treatment of psoriasis, the compounds may be administered orally or topically in amounts sufficient to arrest the proliferation of undifferentiated keratinocytes, and in the treatment of hyperparathyroidism, the compounds should be administered in dosages sufficient to suppress parathyroid activity, so as to achieve parathyroid hormone levels in the normal range. Suitable dosage amounts are from 1 to 500 µg of compound per day, such dosages being adjusted, depending on diseases to be treated, its severity and the response or condition of the subject as well-understood in the art. Typically a composition contains, in a single dosage form, from 0.5 µg to 50 µg of the compound.

This invention is more specifically described by the following illustrative examples. In these examples specific products identified by Roman numerals and letters, i.e. Ia, Ib, ..., IIa, IIb, ..., etc. refer to the specific structures and side chain combinations identified in the preceding description.

Example 1

Preparation of 1α,25-dihydroxy-19-nor-vitamin D₃ (Ia)

(a) 1α,25-Dihydroxy-3,5-cyclovitamin D₃ 1-acetate, 6-methyl etherb: Using 25-hydroxyvitamin D₃ (IIa) as starting material, the known 1α,25-dihydroxy-3,5-cyclovitamin D₃ derivative IIIa (X=H) was prepared according to published procedures (DeLuca et al., U.S. Patent 4, 195,027 and Paaren et al., J. Org. Chem. 45, 3252 (1980)). This product was then acetylated under standard conditions to obtain the corresponding 1-acetate derivative IIIa (X=Ac).

(b) 10,19-Dihydro-1α,10,19,25-tetrahydroxy-3,5-cyclovitamin D₃ 1-acetate, 6-methyl ether (IVa): Intermediate IIIa (X=Ac) was treated with a slight molar excess of osmium tetroxide in pyridine according to the general procedure described by Paaren et al. (J. Org. Chem. 48, 3819 (1983)) to obtain the 10,19-dihydroxylated derivative IVa. Mass spectrum m/z (relative intensity), 506 (M⁺, 1), 488 (2), 474 (40), 425 (45), 396 (15), 285 (5), 229 (30), 133 (45), 59 (80), 43 (100). ¹H NMR (CDCl₃) δ 0.52 (3H, s, 18-CH₃), 0.58 (1H, m, 3-H), 0.93 (3H, d, J=6.1 Hz, 21-CH₃), 1.22 (6H, s, 26-CH₃ and 27-CH₃), 2.10 (3H, s, COCH₃), 3.25 (3H, s, 6-OCH₃), 3.63 (2H, m, 19-CH₂), 4.60 (1H, d, J=9.2 Hz, 6-H), 4.63 (1H, dd, 1β-H), 4.78 (1H, d, J=9.2 Hz, 7-H).

(c) 1α,25-Dihydroxy-10-oxo-3,5-cyclo-19-nor-vitamin D₃ 1-acetate, 6-methyl ether (Va): The 10,19-dihydroxylated intermediate IVa was treated with a solution of sodium metaperiodate according to the procedure given by Paaren et al. (J. Org. Chem. 48, 3819, 1983) to produce the 10-oxo-cyclovitamin D derivative (Va, X=Ac). Mass spectrum m/z (relative intensity) 442 (M⁺-MeOH) (18), 424 (8), 382 (15), 364 (35), 253 (55), 225 (25), 197 (53), 155 (85), 137 (100). ¹H NMR (CDCl₃) δ 0.58 (3H, s, 18-CH₃), 0.93 (3H, d, J=6.6 Hz, 21-CH₃), 1.22 (6H, s, 26-CH₃ and 27-CH₃), 2.15 (s, 3-OCOCH₃), 3.30 (3H, s, 6-OCH₃), 4.61 (1H, d, J=9.1 Hz, 6-H), 4.71 (1H, d, J=9.6 Hz, 7-H), 5.18 (1H, m, 1β-H).

It has been found also that this diol cleavage reaction does not require elevated temperatures, and it is, indeed, generally preferable to conduct the reaction at approximately room temperature.

(d) 1α-Acetoxy-10,25-dihydroxy-3,5-cyclo-19-nor-vitamin D₃ 6-methyl ether (VIa, X=Ac, Y=OH): The 10-oxo derivative Va (X=Ac) (2.2 mg, 4.6 µmol) was dissolved in 0.5 ml of ethanol and to this solution 50 µl (5.3 µmol) of a NaBH₄ solution (prepared from 20 mg of NaBH₄, 4.5 ml water and 0.5 ml of 0.01 N NaOH solution) was added and the mixture stirred at 0°C for ca. 1.5 h, and then kept at 0°C for 16 h. To the mixture ether was added and the organic phase washed with brine, dried over MgSO₄, filtered and evaporated. The crude product was purified by column chromatography on a 15 x 1 cm silica gel column and the alcohol VIa (X=Ac, Y=OH) was eluted with ethyl acetate hexane mixtures to give 1.4 mg (3 µmol) of product. Mass spectrum m/z (relative intensity) 476 (M⁺) (1), 444 (85), 426 (18), 384 (30), 366 (48), 351 (21), 255 (35), 237 (48), 199 (100), 139 (51), 59 (58).

(e) 1α,25-Dihydroxy-19-nor-vitamin D₃ (Ia, X¹=X²=H): The 10-alcohol (VIa, X=Ac, Y=OH) (1.4 mg) was dissolved in 100 µl anhydrous CH₂Cl₂ and 10 µl (14 µmol) triethylamine solution [prepared from 12 mg (16 µl) triethylamine in 100 µl anhydrous CH₂Cl₂], followed by 7 µl (5.6 µmol) mesyl chloride solution (9 mg mesyl chloride, 6.1 µl, in 100 µl anhydrous CH₂Cl₂) added at 0°C. The mixture was stirred at 0°C for 2 h. The solvents were removed with a stream of argon and the residue (comprising compound VIa, X=Ac, Y=CH₃SO₂O-) dissolved in 0.5 ml of anhydrous tetrahydrofuran; 5 mg of LiAlH₄ was added at 0°C and the mixture kept at 0°C for 16 h. Excess LiAlH₄ was decomposed with wet ether, the ether phase was washed with water and dried over MgSO₄, filtered and evaporated to give the 19-nor product VIa (X=Y=H).

This product was dissolved in 0.5 ml of acetic acid and stirred at 55°C for 20 min. The mixture was cooled,

ice water added and extracted with ether. The other phase was washed with cold 10% sodium bicarbonate solution, brine, dried over MgSO_4 , filtered and evaporated to give the expected mixture of 3-acetoxy-1 α -hydroxy- and 1 α -acetoxy-3-hydroxy isomers, which were separated and purified by HPLC (Zorbax SII column, 6.4 x 25 cm, 2-propanol in hexane) to give about 70 μg each of compounds VIIa and VIIIa. UV (in EtOH) λ_{max} 242.5 (OD 0.72), 251.5 (OD 0.86), 260 (OD 0.57).

Both 19-nor-1,25-dihydroxyvitamin D_3 acetates VIIa and VIIIa were hydrolyzed in the same manner. Each of the monoacetates was dissolved in 0.5 ml of ether and 0.5 ml 0.1 N KOH in methanol was added. The mixture was stirred under argon atmosphere for 2 h. More ether was added and the organic phase washed with brine, dried over anhydrous MgSO_4 , filtered and evaporated. The residue was dissolved in a 1:1 mixture of 2-propanol and hexane and passed through a Sep Pak column and washed with the same solvent. The solvents were evaporated and the residue purified by HPLC (Zorbax SII, 6.4 x 25 cm, 10% 2-propanol in hexane). The hydrolysis products of VIIa and VIIIa were identical and gave 66 μg of Ia ($\text{X}^1=\text{X}^2=\text{H}$). Mass spectrum (m/z relative intensity) 404 (M^+) (100), 388 (41), 371 (20), 275 (53), 245 (51), 180 (43), 135 (72), 133 (72), 95 (82), 59 (18), exact mass calcd. for $\text{C}_{26}\text{H}_{44}\text{O}_3$ 404.3290, found 404.3272. ^1H NMR (CDCl_3) δ 0.52 (3H, s, 18- CH_3), 0.92 (3H, d, $\text{J}=6.9$ Hz, 21- CH_3), 1.21 (6H, s, 26- CH_3 and 27- CH_3), 4.02 (1H, m, 3 α -H), 4.08 (1H, m, 1 β -H), 5.83 (1H, d, $\text{J}=11.6$ Hz, 7-H), 6.29 (1H, d, $\text{J}=10.7$ Hz, 6-H). UV (in EtOH), λ_{max} 243 (OD 0.725), 251.5 (OD 0.823), 261 (OD 0.598).

Example 2

Preparation of 1 α -hydroxy-19-nor-vitamin D_3 (Ib)

(a) With vitamin D_3 (IIb) as starting material, and utilizing the conditions of Example 1a, there is obtained known 1 α -hydroxy-3,5-cyclovitamin D_3 1-acetate, 6-methyl ether, compound IIIb ($\text{X}=\text{Ac}$).

(b) By subjecting intermediate IIIb ($\text{X}=\text{Ac}$), as obtained in Example 2a above to the conditions of Example 1b, there is obtained 10,19-dihydro-1 α ,10,19-trihydroxy-3,5-cyclovitamin D_3 1-acetate, 6-methyl ether IVb ($\text{X}=\text{Ac}$).

(c) By treatment of intermediate IVb ($\text{X}=\text{Ac}$) with sodium metaperiodate according to Example 1c above, there is obtained 1 α -hydroxy-10-oxo-3,5-cyclo-19-nor-vitamin D_3 1-acetate, 6-methyl ether Vb ($\text{X}=\text{Ac}$).

(d) Upon reduction of the 10-oxo-intermediate Vb ($\text{X}=\text{Ac}$) under the conditions of Example 1d above, there is obtained 1 α -acetoxy-10-hydroxy-3,5-cyclo-19-nor-vitamin D_3 6-methyl ether VIb ($\text{X}=\text{Ac}$, $\text{Y}=\text{OH}$).

(e) Upon processing intermediate VIb ($\text{X}=\text{Ac}$, $\text{Y}=\text{OH}$) through the procedure given in Example 1e above, there is obtained 1 α -hydroxy-19-nor-vitamin D_3 (Ib, $\text{X}^1=\text{X}^2=\text{H}$).

Example 3

Preparation of 1 α ,25-dihydroxy-19-nor-vitamin D_2

(a) Utilizing 25-hydroxyvitamin D_2 (IIc) as starting material and experimental conditions analogous to those of Example 1a, there is obtained 1 α ,25-dihydroxy-3,5-cyclovitamin D_2 1-acetate, 6-methyl ether, compound IIIc ($\text{X}=\text{Ac}$).

(b) Subjecting intermediate IIIc ($\text{X}=\text{Ac}$), as obtained in Example 3a above, to the reaction conditions of Example 1b, provides 10,19-dihydro-1 α ,10,19,25-tetrahydroxy-3,5-cyclovitamin D_2 1-acetate, 6-methyl ether, IVc ($\text{X}=\text{Ac}$).

(c) By treatment of intermediate IVc ($\text{X}=\text{Ac}$) with sodium metaperiodate according to general procedures of Example 1c above, there is obtained 1 α ,25-dihydroxy-10-oxo-3,5-cyclo-19-nor-vitamin D_2 1-acetate, 6-methyl ether Vc ($\text{X}=\text{Ac}$).

(d) Upon reduction of the 10-oxo-intermediate Vc ($\text{X}=\text{Ac}$) under conditions analogous to those of Example 1d above, there is obtained 1 α -acetoxy-10,25-dihydroxy-3,5-cyclo-19-nor-vitamin D_2 6-methyl ether VIc ($\text{X}=\text{Ac}$, $\text{Y}=\text{OH}$).

(e) Upon processing intermediate VIc ($\text{X}=\text{Ac}$, $\text{Y}=\text{OH}$) through the procedural steps given in Example 1e above, there is obtained 1 α ,25-dihydroxy-19-nor-vitamin D_2 (Ic, $\text{X}^1=\text{X}^2=\text{H}$).

Example 4

Preparation of 1 α -hydroxy-19-nor-vitamin D_2

(a) With vitamin D_2 (IIc) as starting material, and utilizing the conditions of Example 1a, there is obtained known 1 α -hydroxy-3,5-cyclovitamin D_2 1-acetate, 6-methyl ether, compound IIIc ($\text{X}=\text{Ac}$).

(b) By subjecting intermediate IIIc ($\text{X}=\text{Ac}$), as obtained in Example 4a above to the conditions of Example 1b, there is obtained 10,19-dihydro-1 α ,10,19-trihydroxy-3,5-cyclovitamin D_2 1-acetate, 6-methyl ether,

IVd (X=Ac).


(c) By treatment of intermediate IVb (X=Ac) with sodium metaperiodate according to Example 1c above, there is obtained 1 α -hydroxy-10-oxo-3,5-cyclo-19-nor-vitamin D₂ 1-acetate, 6-methyl ether, Vd (X=Ac).

(d) Upon reduction of the 10-oxo-intermediate Vd (X=Ac) under the conditions of Example 1d above, there is obtained 1 α -acetoxy-10-hydroxy-3,5-cyclo-19-nor-vitamin D₂ 6-methyl ether, Vld (X=Ac, Y=OH).

(e) Upon processing intermediate Vld (X=Ac, Y=OH) through the procedure given in Example 1e above, there is obtained 1 α -hydroxy-19-nor-vitamin D₂ (Id, X'=X²=H).


Claims

1. A compound having the formula:



where X¹ and X² are each independently hydrogen, aliphatic acyl of 1 to 5 carbon atoms, aromatic acyl or alkylsilyl or alkoxyalkyl wherein said "alkyl" has 1 to 5 carbon atoms, and R is a side chain known for vitamin D-type compounds:


2. A compound according to claim 1 wherein R is hydroxyalkyl of 1 to 5 carbon atoms, fluoroalkyl of 1 to 5 carbon atoms or a side chain of the formula:



wherein R¹ represents hydrogen, hydroxy or O-acyl, R² and R³ are each independently alkyl, hydroxyalkyl or fluoroalkyl, or, when taken together represent the group $-(CH_2)_m-$ where m is an integer from 2 to 5, R⁴ is hydrogen, hydroxy, fluorine, O-acyl, alkyl, hydroxyalkyl or fluoroalkyl, R⁵ is hydrogen, fluorine, alkyl, hydroxyalkyl or fluoroalkyl, or R⁴ and R⁵ taken together represent double-bonded oxygen, R⁶ and R⁷ are each independently hydrogen, hydroxy, O-acyl, fluorine or alkyl, or, R⁶ and R⁷ taken together form a carbon-carbon double bond, and n is an integer from 1 to 5 and wherein the carbon at any one of positions 20, 22 or 23 in the side chain may be replaced by an O, S, or N atom and wherein said "alkyl" has 1 to 5 carbon atoms and said "acyl" is aliphatic acyl of 1 to 5 carbon atoms or aromatic acyl.


3. A compound according to claim 1 wherein X¹ and X² are both hydrogen, R¹ is hydroxy, R² and R³ are each independently methyl, trifluoromethyl, ethyl or propyl, R⁶ and R⁷ are both hydrogen, or together form a carbon-carbon double bond, R⁴ and R⁵ are both hydrogen and n is 1, 2 or 3.
4. 1 α ,25-Dihydroxy-19-nor-vitamin D₃.

5. 1 α -Hydroxy-19-nor-vitamin D₃.
6. 1 α ,25-Dihydroxy-19-nor-vitamin D₂.
7. 1 α -Hydroxy-19-nor-vitamin D₂.
8. 1 α -Hydroxy-19-nor-24 epi-vitamin D₂.
9. 1 α ,25-Dihydroxy-19-nor-24 epi-vitamin D₂.
10. A compound having the formula:




wherein R is as defined in Claim 1, Q represents alkyl and X is hydrogen, acyl, alkylsilyl or alkoxyalkyl, wherein said "alkyl" has 1 to 5 carbon atoms and said "acyl" is aliphatic acyl of 1 to 5 carbon atoms or aromatic acyl.

11. A compound having the formula:




wherein R is as defined in Claim 1, Q represents alkyl of 1 to 5 carbon atoms and X is as defined in Claim 10.


12. A compound having the formula:



- wherein R is as defined in Claim 1, Q and X are as defined in claim 10, and Y is hydroxy, hydrogen or protected hydroxy where the protecting group is acyl, alkylsilyl or alkoxyalkyl, wherein said "alkyl" has 1 to 5 carbon atoms and said "acyl" is aliphatic acyl of 1 to 5 carbon atoms or aromatic acyl.
13. A pharmaceutical composition which comprises at least one compound as claimed in any one of claims 1 to 9 together with a pharmaceutically acceptable excipient.
 14. A composition according to claim 13 wherein the compound is in a solid or liquid vehicle ingestible by, and non-toxic to, mammals.
 15. A composition according to claim 13 or 14 wherein the compound is 1 α ,25-hydroxy-19-nor-vitamin D₃, 1 α -hydroxy-19-nor-vitamin D₃, 1 α ,25-dihydroxy-19-nor-vitamin D₂ or 1 α -hydroxy-19-nor-vitamin D₂.
 16. A composition according to any one of claims 13 to 15 which contains, in a single dosage form, from 0.5 μ g to 50 μ g of the compound.
 17. A composition according to any one of claims 13 to 16 which is suitable for topical administration.
 18. A composition according to any one of claims 13 to 16 which is suitable for parenteral administration.
 19. A composition according to any one of claims 13 to 16 which is suitable for oral administration.
 20. A compound as defined in any one of claims 1 to 9 for inducing cell differentiation in malignant cells.
 21. A compound as defined in any one of claims 1 to 9 for inducing cell differentiation in leukemia cells.
 22. A compound as defined in any one of claims 1 to 9 for treating a proliferative skin disorder in a mammal.
 23. A compound as defined in any one claims 1 to 9 for treating psoriasis.
 24. A compound as defined in any one of claims 1 to 9 for treating primary or secondary hyperparathyroidism.
 25. A compound as defined in any one of claims 1 to 9 for treating a neoplastic disease.
 26. A process for preparing a compound having the formula:




where X¹, X² and R are as defined in any one of Claims 1 to 3 which comprises solvolysing the 1α-hydroxy-10-deoxy cyclovitamin D compound having the formula:



wherein Q is alkyl of 1 to 5 carbon atoms.

Patentansprüche

1. Verbindung der Formel



worin X¹ und X² unabhängig voneinander Wasserstoff, aliphatisches Acyl mit 1 bis 5 Kohlenstoffatomen, aromatisches Acyl oder Alkylsilyl oder Alkoxyalkyl, worin "Alkyl" 1 bis 5 Kohlenstoffatome besitzt, bedeuten, und R eine für Verbindungen des Vitamin-D-Typs bekannte Seitenkette ist.

2. Verbindung nach Anspruch 1, dadurch gekennzeichnet, daß R Hydroxyalkyl mit 1 bis 5 Kohlenstoffatomen, Fluoralkyl mit 1 bis 5 Kohlenstoffatomen oder eine Seitenkette der Formel ist: